

PURINA® ACCURATION® BALANCER® 26
SUPPLEMENT FOR FEEDLOT CATTLE

What It Is...

A supplement for mixing with coarse cracked corn or shelled corn and by-product ingredients for finishing cattle weighing over 650 lb or dairy beef steers weighing over 400 lb that is formulated to provide intake modifying properties to the complete ration.



Available As:

Product Code	Options	Bag/ Bulk	Size Diameter	Feeding Rate
0040069	R267 T110	Bag	5/32-11/64	1.1-1.6
0040070	R267 T110	Bulk	5/32-11/64	1.1-1.6
0040071	R267 T110	Tote	5/32-11/64	1.25-1.6
0040068	R267 T110 MA 0.2	Bulk	5/32-11/64	1.25-1.6
0040065	R267 T110 MA 0.25	Bag	5/32-11/64	1.25-1.6
0040067	R267 T110 MA 0.25	Bulk	5/32-11/64	1.25-1.6
0040072	B300 AU350	Bulk	5/32-11/64	2
0040073	B300 AU350	Tote	5/32-11/64	2

Features	Benefits
Contains protein, vitamins and minerals	Balances nutrient deficiencies in corn and by product-based ration to improve gains and feed efficiency
Formulated using proprietary intake modifying technology	Regulates number and size of meals which improves feed efficiency and reduces digestive and metabolic problems in finishing cattle including bloat, acidosis and founder. Reduces need for roughage in ration which improves feed efficiency, reduces storage and handling costs as well as decreasing manure production
Drug options available	Increase gains and feed efficiency

PURINA® ACCURATION® BALANCER® 26 R267 T110

SUPPLEMENT FOR FEEDLOT CATTLE

For improved feed efficiency. For reduction of incidence of liver abscesses in beef cattle caused by fusobacterium necrophorum and corynebacterium.

ACTIVE DRUG INGREDIENTS:

Monensin (as Monensin Sodium).....	267.0 G/T
Tylosin (as Tylosin Phosphate).....	110.0 G/T

CAUTION: USE ONLY AS DIRECTED
INTENDED FOR FEEDING BEEF CATTLE ONLY

GUARANTEED ANALYSIS

Crude Protein (min.).....	26.0%
This includes no more than 17% equivalent crude protein from non-protein nitrogen	
Crude Fat (min.).....	2.0%
Crude Fiber (max.).....	15.0%
Calcium (Ca) (min.).....	6.0%
Calcium (Ca), max.	7.0%
Phosphorus (P) (min.).....	0.5%
Salt (NaCl) (min.).....	2.5%
Salt (NaCl) (max.).....	3.5%
Potassium (K) (min.).....	2.0%
Vitamin A (min.).....	20,000 IU/lb

MIXING DIRECTIONS:

Thoroughly mix 170 pounds of this product with 1830 pounds of good quality air dry coarse cracked or whole shelled corn and/or byproducts to yield a final mixed ration containing 9.35 grams/ton tylosin and 22.7 grams/ton monensin.

FEEDING DIRECTIONS

Feed this final mix continuously as the sole ration to finishing beef cattle weighing 650 pounds or more or dairybeef steers weighing 400 pounds or more at the rate of 19.25 pounds per head per day to provide 218 mg. monensin and 90 mg. tylosin per head per day. No additional improvement in feed efficiency has been shown from feeding monensin at levels greater than 30 g/ton (360 mg monensin per head per day).

LIMITATIONS FOR USE:

Feed only to cattle being fed in confinement for slaughter.

IMPORTANT:

This product is to be fed only to the animal species as directed on this label. Follow these management practices:

1. Cracked corn should be processed in a manner to minimize the creation of fines.
2. Whole shelled corn should be at least of #2 corn quality.
3. Storage and handling of supplement should be managed to eliminate the unnecessary production of fines.
4. Do not allow complete ration fines to build up in the feed bunk or bulk feeder.
5. Proper adaptation procedures to the complete ration should be followed.
6. When making a ration change, allow 7-10 days for animals to adjust to the new ration.
7. Provide a source of fresh, clean water at all times.
8. Provide adequate bunk space for each animal. Bunks should be well protected and well managed to prevent feed from becoming wet and molding.
9. When fed from a self-feeder, adjust feeder to minimize quantity of feed accumulating in trough.
10. Cattle should be vaccinated against enterotoxemia.
11. Consult your veterinarian for recommended health program for your local area.
12. Have feed before cattle at all times.